

Source:
Dr. David Michael Glenn USDA, ARS, West
Virginia, USA


12% Surround

Control

3% Surround




Source:
Dr. David Michael Glenn USDA, ARS, West
Virginia, USA




In summer the use particle film (kaolin particles) applied in high light condition (clear sky)

- Increase reflection of UV and IR
- Reduce leaf temperature
- Increase stomatal conductance
- Increase whole-canopy photosynthesis and transpiration


Concentration: 10%/12%



12kg = 38US\$




Effects on sap flow




non-heated probe

Sap flow measure
differences
between
heated and
non-heated probe



Water reduce
temperature



May to July (winter dry season)(104days)

Plant leaf area: **5m²**

Kaolin particles:

$$0.70 L h m^{-2} \times 5m^2 = 3.5 L h^{-1} plant^{-1} \times 8h = \mathbf{28 L H_2O plant^{-1} day^{-1}}$$

Control:

$$0.32 L h m^{-2} = 1.60 L h plant \times 8h = \mathbf{12.8 L H_2O plant^{-1} day^{-1}}$$

$$\mathbf{Maximum light = 2300\mu mol m^{-2} s^{-1} = 1000 W m^{-2}}$$



I think that Kaolin particles is important to papaya:

- Can reduce midday photosynthesis depression (stomatal effects is so important to increase yield in papaya). The photosynthetic capacity of a papaya genotype also influences papaya fruit quality (Salazar, 1978).
- Can control aphids ([Aphids](#) are the predominant means by which papaya ring spot virus (PRSV) is transmitted)
- Reduce canopy temperature (Prolonged droughts, associated with high temperatures, adversely affect fruit production because the condition induces abortion of floral and fruit structures, leading to sterile phases or fruiting skips along the stem)
- Increase scatter light inside the greenhouse
- Kaolin particles reduce dew formation on leaves

Salazar, R. 1978. Determination of photosynthesis in commercial varieties of papaw (*Carica papaya* L.) and its possible relationship with the production and quality of the fruits. Revista ICA. Bogota 13(2):291-295.





Fig. 1. Net CO_2 assimilation (A) of leaves from control (■) and papaya ringspot virus-infected (▲) *Carica papaya* plants as a function of incident photosynthetic photon flux (PPF). Each point represents the mean of six replications (as determined on 5–6 Jan. 1992). Equations for the response curves are $A = (-21.87) + 5.32 \ln \text{PPF}$, $r^2 = 0.98$ (control); and $A = (-15.77) + 3.37 \ln \text{PPF}$, $r^2 = 0.98$ (diseased); n was smaller than symbols where no bars are visible.



Table 1. Influence of papaya ringspot virus on net CO_2 assimilation at saturating light and ambient CO_2 (A_{max}), internal CO_2 partial pressure at A_{max} (C_{max}), CO_2 use efficiency, apparent quantum yield (ϕ), the ratio of variable to maximal fluorescence (F_v/F_{max}), and dark respiration (R_d) of 'Kapoho' papaya plants. Measurements were made on systemically infected leaves, 7 to 8 weeks after inoculation ($n = 7$ for F_v and 6 for others).

Variable	Control	Diseased
A_{max} ($\mu\text{mol CO}_2 / \text{m}^2 \text{ per sec}$)	15.7	8.9**
C_{max} ($\mu\text{bar CO}_2$)	273	267*
CO_2 use efficiency ($\mu\text{mol CO}_2 / \mu\text{bar CO}_2$)	0.053	0.030***
ϕ ($\mu\text{mol CO}_2 / \mu\text{mol photons}$)	0.039	0.016***
F_v/F_{max}	0.80	0.76*
R_d ($\mu\text{mol CO}_2 / \text{m}^2 \text{ per sec}$)	1.79	2.94**

* ** *** Nonsignificant or significant at $P = 0.05$, 0.01, or 0.001, respectively.



Symptoms of Papaya Ringspot Virus
(photo credit: John McHugh)

Action of Particle Films Against Insects and Mites



1. Repellent
2. Oviposition Deterrent
3. Feeding Inhibitor
4. Acute Mortality
5. Chronic Mortality

6. Impedes Grasping (Fall-Off)
7. Paralysis or Altered Behavior
8. Host Camouflaging
9. Restricts Movement or Infestation Process



Inside the greenhouse light is very
limitant









Kaolin particles reduce dew in papaya leaves

Can reduce disease!

Microspray irrigation applying above the canopy during summer (12h to 14h).
Evaporative cooling reduce $VPD_{\text{leaf-air}}$, increase stomatal conductance, increase transpiration and photosynthesis



When the air temperature inside the canopy reached **35°C** the microspray irrigation system was turned on



Microspray
irrigation
increased

5 fruits plant⁻¹



Greenhouse can provides adequate environmental conditions to cultivate papaya.

- protection against excessive light (sunburn, photoinhibition, high leaf temperature, high $VPD_{\text{leaf-air}}$)
- protection against wind and hail
- Exclusion of PRV vector







Baixinho de Santa Amália genotype












30% Light reduction










Copyright © 2005 Pearson Education, Inc. Publishing as Pearson Benjamin Cummings. All rights reserved.

Commercial production

2004/2005	Greenhouse	Full sunlight
Plant Height (cm)	183.8 a	144.2 b
Trunk diameter (cm)	13 a	10 b
Leaf number plant ⁻¹	35.3 a	29.4 b
Leaf area plant (m ²)	0.2 a	0.15 b
Number of fruits plant ⁻¹	9.7 a	6.5 b
Fruit weight plant ⁻¹	3.53 kg a	2.12 kg b
Average fruit weight	364.7 g a	326.1g b
Martelleto et al 2008		

Light quality and intensity can affect leaf anatomy in papaya plant





< R/FR



Shade 50%



Full sunlight Buisson e Lee, 1993



Buisson e Lee, 1993

	Sun leaf	Shade leaf
Leaf thickness (μm)	137	119
Specific leaf mass (mg cm^{-2})	4,7	2,8
Leaf area (cm^2)	292	162
Chlorophyll content ($\mu\text{g cm}^{-2}$)	3.57	5.16
Petiole lenght (mm)	207	148
Stomata density (mm^{-2})	465	330
Degree air space	0.29	0.33

Under conditions of low light intensity and low red:far red light , leaf lobing was dramatically reduced.


Change in spectral quality also resulted in a reduction in the ratio chlorophyll a to b



High Leaf lobing can reduce leaf temperature in sun leaf



Motorcycle radiator




The papaya photosynthetic capacity can be linked to non stomatal limitation in soil field capacity condition.

As exemple:

-Nitrogen leaf concentration

15 g kg⁻¹DM

35 g kg⁻¹DM



18 g kg⁻¹DM

37 g kg⁻¹DM






Fig. 2 Relationship between chlorophyll concentration (a+b) ($\mu\text{mol m}^{-2}$) and leaf N_{org} (g kg^{-1} DW) in 'Solo' and 'Golden' papaya plants cultivated in greenhouse conditions. The R^2 indicated significant correlation at the 0.1 % level










Figure 4. The relation between the organic nitrogen content and the PCM reading, in Sunrise Solo and Golden papaya plants, cultivated in a greenhouse. The open circles with reduced PCM readings (<30) refer the senescent 6th – 7th leaves.








Fig. 6 Relationship between total maximum quantum efficiency (F_v/F_m) and SPAD values in 'Solo' and 'Golden' papaya plants cultivated in greenhouse conditions. R^2 indicate significant correlation at 0.1 % level




Fig. 4 Relationship between chlorophyll concentration (a + b) ($\mu\text{mol m}^{-2}$) and SPAD values in 'Solo' and 'Golden' papaya plants cultivated in greenhouse conditions. R^2 indicate significant correlation at the 0.1 % level




Fig. 3 Relationship between net photosynthetic rate (A) ($\mu\text{mol m}^{-2} \text{s}^{-1}$) and chlorophyll concentration ($a + b$) ($\mu\text{mol m}^{-2}$) in 'Solo' and 'Golden' papaya plants cultivated in greenhouse conditions. R^2 indicate significant correlation at the 0.1 % level

Portable chlorophyll meter (PCM-502) values are related to total chlorophyll concentration and photosynthetic capacity in papaya (*Carica papaya* L.)

Fernanda Assumpção De Castro · Elienar Campestre · Alena Torres Netto ·
Mara De Menezes De Azeite Gomes · Thiago Manoel Ferraz · David Michael Glenn







Figure 3. Relationship between the RC/CS_a ratio (density of active reaction center related to reduction in Qa) and the PCM reading, in Sunrise Solo and Golden papaya plants cultivated in a greenhouse. The open circles with reduced PCM readings (<30) refer the senescent 6th – 7th leaves.



Relationship between photochemical efficiency (JIP-Test Parameters) and portable chlorophyll meter readings in papaya plants

Fernando Assumpção Castro¹, Eliemar Campestri^{1*}, Alena Torres Netto¹,
Leandro Hespanhol Viana¹

1 Spad reading



45 Spad reading








Figure 8. Energy pipeline models of the specific (membrane model, per RC) and phenomenological (leaf model, per CS) fluxes of leaf from a papaya plant of the Sunrise Solo genotype with PCM values of 5.0 (A) and 45 (B), and the Golden genotype with PCM values of 1.0 (C) and 39 (D). The solid and open circles show inactive and active RCs, respectively. The value of each of the parameters can be seen in the relative changes in the width of each arrow.







Fig. 1 Relationship between net photosynthetic rate (A) ($\mu\text{mol m}^{-2} \text{s}^{-1}$) and leaf N_{org} ($\text{g kg}^{-1} \text{DW}$) in 'Solo' and 'Golden' papaya plants cultivated in greenhouse conditions. The R^2 indicated significant correlation at the 0.1 % level. Fig. (b) is the linear part of the Fig. (a) and is related to the photosynthetic nitrogen use efficiency





Fig. 2 Relationship between chlorophyll concentration (a+b) ($\mu\text{mol m}^{-2}$) and leaf N_{org} ($\text{g kg}^{-1} \text{DW}$) in 'Solo' and 'Golden' papaya plants cultivated in greenhouse conditions. The R^2 indicated significant correlation at the 0.1 % level

positively correlated with chlorophyll content (Fig. 2) although photosynthesis was curvilinearly related to

Blade Leaf: Optimal N concentration **50 g Kg^{-1}** (Marinho 2002)

Petiole: Optimal N concentration **11 g Kg^{-1}** (Caliman Company, Brazil)




Fig. 5 Relationship between photosynthesis (A) ($\mu\text{mol m}^{-2} \text{s}^{-1}$) and SPAD values in 'Solo' and 'Golden' papaya plants cultivated in greenhouse conditions. R^2 indicate significant correlation at the 0.1 % level

Soil and air water

1) Air water:



Threshold VPD values to papaya:

$$VPD_{\text{air}} = < 1\text{kPa} (e_{\text{sair}} - e_{\text{air}})$$

$$VPD_{\text{leaf-air}} = < 2\text{kPa} (e_{\text{sleaf}} - e_{\text{air}})$$

2) Soil water:

Papaya exhibits both stomatal and non stomatal response to soil water deficits and the source of the response signals are both hydraulic and non-hydraulic in nature



Responses of papaya seedlings (*Carica papaya* L.) to water stress and re-hydration: growth, photosynthesis and mineral nutrient imbalance

Jakel Mahowach^{1,3}, Ana Rosa Socorro¹ & Manuel Talon²

¹Departamento de Suelos y Riegos, Instituto Canario de Investigaciones Agrarias, Apdo 60, E-38200 La Laguna, Santa Cruz de Tenerife, Spain, ²Centro de Genómica, Instituto Valenciano de Investigaciones Agrarias, E-46113 Moncada, Valencia, Spain, ³Corresponding author*




Figure 2. Moisture (%) in watered (○) and non-watered pot soils (●) of papaya seedlings. In non-watered soils, irrigation was suspended during 34 days and re-established thereafter until the end of experiment. Data are means $\pm$ SE and each value was determined by three TDR probes with three replicates per treatment ($n \geq 9$) (one probe per pot). DAT = days after treatment. WS: Water Stress. RH: Re-hydration.




Figure 1. Leaf water potential in papaya seedlings. Treatments were: regular irrigated plants (○) and non-irrigated plants during 34 days followed by re-watering until the end of experiment (●). Data are means $\pm$ SE and each value was determined in at least three different plants with three replicates per treatment ($n \geq 9$). DAT = days after treatment. WS: Water Stress. RH: Re-hydration.

Baixinho de Santa Amália genotype
70L pots
Greenhouse
Maximum PAR $1200 \mu\text{mol m}^{-2} \text{s}^{-1}$
Severe water stress: $\psi_{\text{leaf}} = -0.8 \text{ MPa}$
Regular irrigated plants: $\psi_{\text{leaf}} = -0.6 \text{ MPa}$

Table 1. Plant height and stem circumference of papaya seedlings subjected to water stress and re-hydration. The treatment consisted of irrigation suppress during 34 days and re-watering establishment thereafter until 48 days (total experimental period). The data ($n \geq 9$) presented in each line followed by dissimilar letters differ significantly at ($P \leq 0.05$). *: $P \leq 0.01$. **: $P \leq 0.001$

Treatments	Plant height (cm)		Stem circumference (cm)	
	Control	Water stress	Control	Water stress
Water stress (Days)				
0	$38.45 \pm 0.67a$	$38.50 \pm 0.56a$	$8.23 \pm 0.09a$	$8.08 \pm 0.15a$
14	$41.67 \pm 0.80a$	$39.63 \pm 0.58a$	$9.04 \pm 0.10a$	$8.72 \pm 0.13a$
20	$42.21 \pm 1.06a$	$39.88 \pm 0.60b$	$9.51 \pm 0.11a$	$8.94 \pm 0.15b^*$
34	$43.83 \pm 1.15a$	$40.13 \pm 0.67b$	$9.79 \pm 0.12a$	$8.98 \pm 0.12b^{**}$
Rehydration (Days)				
41	$46.13 \pm 1.28a$	$41.20 \pm 0.87b$	$10.10 \pm 0.23a$	$8.98 \pm 0.19b^*$
48	$46.83 \pm 1.86a$	$43.25 \pm 1.75b$	$10.60 \pm 0.32a$	$9.40 \pm 0.35b$




Figure 3. Relative growth rate using stem DW of papaya seedlings. Treatments were: regular irrigated plants (□) and non-irrigated plants during 34 days followed by re-watering until the end of experiment (▨). Data are means $\pm$ SE and each value was determined in at least three different plants with three replicates per treatment ($n \geq 9$). DAT = days after treatment, WS: Water Stress, RH: Re-hydration.

Table 2. Root fresh weight of papaya seedlings subjected to water stress and re-hydration. The treatment consisted of irrigation suppress during 34 days and re-watering establishment thereafter until 48 days (total experimental period). The data ($n \geq 9$) presented in each line followed by dissimilar letters differ significantly at ($P \leq 0.05$)

Treatments	Root fresh weight (g)	
	Control	Water stress
Water stress (Days)		
34	181.9 $\pm$ 16.5a	101.4 $\pm$ 18.7b
Rehydration (Days)		
48	183.0 $\pm$ 23.5a	119.7 $\pm$ 8.70b




Figure 4. Number of remaining leaves in papaya seedlings. Treatments were: regular irrigated plants (○) and non-irrigated plants during 34 days followed by re-watering until the end of experiment (●). Data are means $\pm$ SE and each value was determined in at least three different plants with three replicates per treatment ($n \geq 9$). DAT=days after treatment. WS: Water Stress. RH: Re-hydration.




Figure 5. Photosynthetic rate (a) in fully expanded leaves in papaya seedlings. Treatments were: regular irrigated plants (○) and non-irrigated plants during 34 days followed by re-watering until the end of experiment (●). Data are means $\pm$ SE and each value was determined in at least three different plants with three replicates per treatment ($n \geq 9$). DAT=days after treatment. WS: Water Stress. RH: Re-hydration.




Figure 7. Potassium content in leaf (A) and root (B) tissues in papaya seedlings. Treatments were: regular irrigated plants (○) and non-irrigated plants during 34 days followed by re-watering until the end of experiment (●). Each value is the mean of at least three independent measurements ($n \geq 3$) $\pm$ SE. DAT=days after treatment, WS: Water Stress, RH: Re-hydration.




Figure 8. Sodium content in leaf (A) and root (B) tissues in papaya seedlings. Treatments were: regular irrigated plants (○) and non-irrigated plants during 34 days followed by re-watering until the end of experiment (●). Each value is the mean of at least three independent measurements ($n \geq 3$) $\pm$ SE. DAT=days after treatment, WS: Water Stress, RH: Re-hydration.




Figure 9. Chloride content in leaf (a) and root (b) tissues in papaya seedlings. Treatments were: regular irrigated plants (○) and non-irrigated plants during 34 days followed by re-watering until the end of experiment (●). Each value is the mean of at least three independent measurements ($n \geq 3$) $\pm$ SE. DAT=days after treatment. WS: Water Stress. RH: Re-hydration.

In conclusion, papaya plants subjected to water stress showed a tendency to accumulate ions such as K^+ , Na^+ and Cl^- . The ion increases that were registered on per DW basis might apparently contribute for osmotic adjustment, enhancing water stress tolerance of these plants. The data also indicate that stress tolerance was not mediated through the reduction of leaf abscission, the detention of growth or decrease of net CO_2 assimilation. Re-irrigation induced plant morphological and physiological recovery without irreversible effects of water stress.

Hormonal changes in papaya seedlings subjected to progressive water stress and re-watering

Jalel Mahbouchi · Vicent Arbona ·
 Aurelio Gómez-Cadenas




Fig. 1 Soil moisture (%) in irrigated (□) and non-irrigated pot soils (■) of papaya seedlings. In water-stressed soils, irrigation was released during 40 days and re-established thereafter until the end of experiment. Data are means $\pm$ SE and each value was determined by three TDR probes with three replicates per treatment ($n \geq 9$) (one probe per pot). DAT = days after treatment. WS: water stress, RH: rehydration




Fig. 2 Principal leaf vein length (a) and leaf blade dry weight (b) in papaya seedlings. Treatments were: regular irrigated plants (○) and water-stressed plants during 40 days followed by re-watering until the end of experiment (●). Data are means $\pm$ SE and each value was determined from at least three different plants with three replicates per treatment ($n \geq 9$). DAT = days after treatment. WS: water stress, RH: rehydration




Fig. 3 Percentage of leaves present in papaya plants subjected to WS during 40 days and subsequent RH (■). Data are expressed as percentages with respect to control plants (---) in each date. Data are means $\pm$ SE and each value was determined in at least three different plants with three replicates per treatment ($n \geq 9$). DAT = days after treatment. WS: water stress, RH: rehydration




Fig. 4 Photosynthetic rate (A, (a)), stomatal conductance (gs, (b)) and transpiration rate (E, (c)) in fully expanded leaves in papaya seedlings. Treatments were: regular irrigated plants (○) and water-stressed plants during 40 days followed by re-watering until the end of experiment (●). Data are means $\pm$ SE and each value was determined in at least three different plants with three replicates per treatment ($n \geq 9$). DAT = days after treatment. WS: water stress, RH: rehydration








Fig. 5 Abscisic acid (ABA) levels in leaves (a) and roots (b) from well-watered plants (□) and from plants subjected to WS during 40 days and subsequent RH (■) in papaya seedlings. Each value is the mean of at least three independent measurements ($n \geq 3$) $\pm$ SE. DAT = days after treatment. WS; water stress, RH: rehydration




Field-grown papaya



21 Mar. 1996, when ψ_m was -62 kPa for the drought-stressed obtained on 29 Mar., when ψ_m was -68 kPa for the drought

Treatment	ϕ_i (mol CO ₂ /mol incident quanta)
Control	0.039 ± 0.001
Drought	$0.025 \pm 0.001^{**}$
Control	0.044 ± 0.001
Drought	$0.024 \pm 0.001^{**}$
Control	0.040 ± 0.001
Drought	$0.024 \pm 0.001^{**}$


^{**}Significant at $P = 0.01$.



Marler e Mickelbart, 1998

Strategies to increase effective use of water in papaya

- Regulated deficit irrigation (RDI)
- Partial rootzone drying (PRD)



Partial rootzone drying (PRD) and regulated deficit irrigation (RDI) effects on stomatal conductance, growth, photosynthetic capacity, and water-use efficiency of papaya^a



Roberta Samara Nunes de Lima^a, Fábio Afonso Mazzei Moura de Assis Figueiredo^a, Amanda Oliveira Martins^a, Bruna Corrêa da Silva de Deus^a, Tiago Massi Ferraz^a, Mara de Menezes de Assis Gomes^a, Elias Fernandes de Sousa^b, David Michael Glenn^c, Eliemar Campostrini^{a,*}

^a Setor de Fisiologia Vegetal, EMCV, Centro de Ciências e Tecnologias Agropecuárias, Universidade Estadual do Norte Fluminense, Av. Alberto Carmo, 2000, CEP: 28013-620 Campos dos Goytacazes, RJ, Brazil

^b Setor de Irrigação e Drenagem, LEAG, Centro de Ciências e Tecnologias Agropecuárias, Universidade Estadual do Norte Fluminense, Av. Alberto Carmo, 2000, CEP: 28013-620 Campos dos Goytacazes, RJ, Brazil

^c USDA-ARS, Appalachian Fruit Research Station, 2217 Wilshire Road, Kearneysville, WV 25420, USA

Strategies to increase effective use of water in papaya

- Regulated deficit irrigation (RDI)
- Partial rootzone drying (PRD)

15L pots

Substrate soil, sand and cattle manure (2:1:2)





The plants were kept at field capacity (FC) until they were 96 days old.







- Gas exchange
- Chlorophyll fluorescence
- Growth (central vein length, root dry biomass, stem dry biomass, leaf dry biomass, total dry biomass, root volume)
- Proline
- Carbon isotope discrimination
- Agronomic water use efficiency
- Thermal imaging



The plants were kept at field capacity (FC) until they were 96 days old.

Table 1
Timeline of irrigation treatments.

		Days after irrigation treatment application																					
Treatment	Pot	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
FI	1	100% FC																					
	2	100% FC																					
PRD	1	100% FC															100% FC						
	2									100% FC													
RDI	1											50% FC											
	2											50% FC											
NI	1															100% FC							
	2															100% FC							

Full irrigated (FI); partial-root-zone drying (PRD); regulated deficit irrigation (RDI); non-irrigated (NI). Dark-grey represents the period of days in which irrigation was applied at 100% of field capacity (FC). The light grey represents the days with irrigation at 50% of field capacity. The white color represents the period of time with no irrigation.

Table 1

Total volume of water applied, and volume applied per day of greenhouse-grown papaya (*Carica papaya* L.) in splitroot pots with four different irrigation treatments: full irrigated (FI), Partial Rootzone Drying (PRD), Regulated Deficit Irrigation (RDI), and non-irrigated followed by 6 days of FI (NI-gh).

Treatment	Total volume of water applied (L)	Volume of water applied per day (L)
FI	47.50	2.3
PRD	23.8	1.1
RDI	23.8	1.1
NI-gh	21.3	1.0





Fig. 1. Daily soil water potential (-kPa) planted with 'Grand Golden' papaya with four irrigation treatments. PRD-1 and PRD-2 indicate the two sides of the PRD pots.



NI


PRD

14 days after planting
maximum stress

Fig. 10. Relationship between stomatal conductance and soil water potential in papaya in four irrigation treatments in the greenhouse.



NI









FI

NI



**14 days after planting
maximum stress**











Fig. 6. Relationship of leaf-air vapor pressure deficit to transpiration and stomatal conductance in papaya in four irrigation treatments in a greenhouse.






Fig. 3. Relationship of stomatal conductance (G_s) to transpiration (E), photosynthesis (A) and water use efficiency (A/E) for papaya treated with four irrigation regimes.




Fig. 5. Relationship of photosynthesis to stomatal conductance with the leaf-air vapor pressure deficit in papaya treated with four irrigation treatments in the greenhouse.

Photochemical efficiency of PSII





$$PI = (RC/ABS) \times (TR/DI) \times (ET/(TR-ET))$$

(RC/ABS): Active RC density on a Chl basis

(F_v/F_0): Performance due to trapping probability
 $F_v/F_0 = TR/DI$

($ET/(TR-ET)$): Performance due to electron-transport probability

$$F_v/F_m = TR/ABS$$



Copyright © 2005 Pearson Education, Inc. Publishing as Pearson Benjamin Cummings. All rights reserved.




Fig. 7. Central vein length during the ontogeny of the youngest leaf of each plant ('Grand Golden' papaya) in splitroot pots under four different irrigation regimes: FI, NI-gh, PRD, RDI. The narrow and broad arrows indicate the first alternating between the two root sides of the PRD pots and rehydration of NI treatment, respectively ($n = 10$).







Fig. 11. Relationship between incremental leaf growth (% of full irrigation) and soil water potential in papaya in four irrigation treatments in the greenhouse.








Fig. 12. Relationship of stomatal conductance (G_s) to incremental leaf growth and photosynthesis (A) in papaya in four irrigation treatments in a greenhouse.




Fig. 8. Biomass allocation of leaf, trunk and root ($\text{g dry weight plant}^{-1}$) tissues in papaya treated with four irrigation regimes in the greenhouse. FI, NI-gh, PRD, RDI. Capital letters refer to mean separation between treatments. Lower case letters refer to mean separation within the three tissue components ($P=0.05$).






Fig. 9. Relationship of photosynthesis to dry weight 14 days after initiating treatments (the day of most intense water stress for the plants in NI-gh) in 110-days-old 'Grand Golden' papaya plants grown in splitroot pots under four different irrigation regimes (FI, NI-gh, PRD, RDI) in a greenhouse. Values represent means of 10 replicates.



FI



NI



RDI



PRD



FI



FI




NI




Dry side of PRD

**root dead
(7 days without
irrigation)**

**young root
5 days with
irrigation
after 7 days
without
irrigation**





Carbon isotope discrimination




$$>C_i/C_a (\approx 0,7) = < \delta \text{ ‰}$$

$$<C_i/C_a (\approx 0,3) = > \delta \text{ ‰}$$



Agronomic water use efficiency



Treatment	Water economy in relation FI (%)	AWUE (g DM L ⁻¹)	Transpiration rate (L H ₂ O g DM ⁻¹)
FI	-	3.12	0.322
PRD	50%	4.55	0.217
RDI	50%	4.57	0.217
NI	55.14%	2.46	0.400

C3 crops

1 to 6 g DM L⁻¹ H₂O

C4 grasses

10 to 30 g DM L⁻¹ H₂O

Arkley (1982)

Table 1

Total volume of water applied, and volume applied per day of greenhouse-grown papaya (*Carica papaya* L.) in splitroot pots with four different irrigation treatments: full irrigated (FI), Partial Rootzone Drying (PRD), Regulated Deficit Irrigation (RDI), and non-irrigated followed by 6 days of FI (NI-gh).

Treatment	Total volume of water applied (L)	Volume of water applied per day (L)
FI	47.50	2.3
PRD	23.8	1.1
RDI	23.8	1.1
NI-gh	21.3	1.0

Treatment	L H ₂ O m ⁻² day ⁻¹
FI	1.63
PRD	0.84
RDI	0.78
NI	1.17



Treatment	Volume water applied per plant per day	Transpiration L H2O per m ² leaf per day per plant	Transpiration L H2O per plant per day	Leaf area m ²	age
Whole canopy summer	16.0	2.5	10	4.0	5 months
Whole canopy winter	10.0	4.2	15	3.5	5 months
FI	2.3	1.63	2.3	1.41	3 months
PRD	1.1	0.84	1.1	1.30	3 months
RDI	1.1	0.78	1.1	1.40	3 months
NI	1.0	1.17	1.0	0.85	3 months

Thermal imaging

G Model

AGWAT-4235; No. of Pages 10

ARTICLE IN PRESS

6

R.S.N. Lima et al. / Agricultural Water Management xxx (2015) xxx–xxx




Fig. 3. (A) Leaf temperature (T_{leaf}) derived from IR measurements during the study (DAT); (B) difference of leaf to air temperature ($\Delta T_{leaf-air}$), and (C) false-colored IR thermal images showing a selected fully expanded leaf, along the experiment for the different treatments: fully irrigated (FI); partial root drying (PRD); regulated-deficit irrigation (RDI); non-irrigated (NI). Climate conditions at 0 DAT (T_{air} max: 25 °C, RH_{max} : 80%, ψ_{soil} = -10 kPa, PAR_{max} = 257 $\mu\text{mol m}^{-2} \text{s}^{-1}$); 9 DAT (T_{air} max: 44 °C, RH_{max} : 26%, PAR_{max} = 900 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and 14 DAT (T_{air} max: 36 °C, RH_{max} : 39%, PAR_{max} = 830 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Leaf temperature scale is identical for four treatments in the same day of observation. Different letters indicate significant differences at $P < 0.05$ by the Tukey's test ($n = 10$).

Linking thermal imaging to physiological indicators in *Carica papaya* L. under different watering regimes^a

R.S.N. Lima^{a,*}, I. García-Tejero^{a,c}, T.S. López^d, J.M. Costa^e, M. Vaz^f, V.H. Durán-Zuazo^e,
 M. Chaves^g, D.M. Glenn^h, E. Compañónⁱ




Fig. 4. Relationship between transpiration (E), stomatal conductance to water vapor (g_s) and net assimilation (A_n) vs. the leaf temperature (T_{leaf}) and the difference between T_{leaf} and air temperature ($\Delta T_{leaf-air}$). Each linear function was determined with 80 pair of data ($n = 80$).




Fig. 5. Responses of transpiration (E), stomatal conductance (g_s), and net photosynthesis (A_n) to a temperature step (ΔT_{leaf}).





Contents lists available at ScienceDirect

Scientia Horticulturae

journal homepage: www.elsevier.com/locate/scihorti



Partial rootzone drying (PRD) and regulated deficit irrigation (RDI) effects on stomatal conductance, growth, photosynthetic capacity, and water-use efficiency of papaya^{a,*}

Roberta Samara Nunes de Lima^a, Fábio Afonso Mazzei Moura de Assis Figueiredo^a, Amanda Oliveira Martins^a, Bruna Corrêa da Silva de Deus^a, Tiago Massi Ferraz^a, Mara de Menezes de Assis Gomes^a, Elias Fernandes de Sousa^b, David Michael Glenn^c, Eliemar Campostrini^{a,*}

^a Setor de Fisiologia Vegetal, LIMCZ, Centro de Ciências e Tecnologias Agropecuárias, Universidade Estadual do Norte Fluminense, Av. Alberto Lamego, 2000, CEP: 28091-3620 Campos dos Goytacazes, RJ, Brazil

^b Setor de Irrigação e Drenagem, LEAG, Centro de Ciências e Tecnologias Agropecuárias, Universidade Estadual do Norte Fluminense, Av. Alberto Lamego, 2000, CEP: 28091-3620 Campos dos Goytacazes, RJ, Brazil

^c USDA-ARS, Appalachian Fruit Research Station, 2217 Wiltshire Road, Kearneysville, WV 25430, USA



20 months
Caliman company
Brazil
<http://www.caliman.com.br/pt/>




Field condition



Field condition

ET_0



Treatment ²	Irrigation + precipitations (L)
FI	2698
NI-field	1025
PRD100	2698
PRD70	2189
RDI	2189
FI	3755
NI-field	1107
PRD100	3755
PRD70	2949
RDI	2949

measurement period (Jones, 1992). The meteorological station was installed 100 m in the experiment, and the data were used to calculate the reference evapotranspiration (ET_0) using the Penman equation parameterized by the United Nations Food and Agriculture Organization (FAO) (Pereira et al., 1997) (Eq. (1)). We considered that the daily balance heat flow in soil was zero ($G=0$).

$$ET_0 = \frac{1}{(1 + \gamma^*)} \left(R_n - G \right) \frac{1}{k} + \frac{\gamma^*}{(1 + \gamma^*)} \frac{U_2}{(T + 273.15)} (e_s - e_a) \quad (1)$$

where: s is the slope of the vapor pressure curve ($kPa \cdot ^\circ C^{-1}$); γ^* is the modified psychrometric constant ($kPa \cdot ^\circ C^{-1}$); R_n is the net radiation ($MJ m^{-2} d^{-1}$); G is the heat flow in soil ($MJ m^{-2} d^{-1}$); λ is the latent heat of evaporation ($MJ kg^{-1}$); γ = psychrometric coefficient ($kPa \cdot ^\circ C^{-1}$); T is the average temperature ($^\circ C$); U_2 is the wind speed at 2 m ($m s^{-1}$); e_s is the saturation vapor pressure (kPa); e_a is the partial pressure of vapor (kPa).




Fig. 2. Meteorological variables in a field study.

to 5-month-old plants: [redacted] was applied on both sides of the root with 1 drip line and 2 emitters per plant (0.75 m emitter spacing). Emitter flow was $2.3 L h^{-1}$ (total flow: $4.6 L plant^{-1}$); [redacted] was applied to one side only of the root, and every 15 days, water was applied to the alternate side of the root system. Water was applied with 2 drip lines and 2 emitters per plant (0.75 m emitter spacing). Emitter flow was $2.3 L h^{-1}$ (total flow: $4.6 L plant^{-1}$); [redacted] was applied to one side only of the root, and every 15 days, water was applied to the alternate side of the root system to allow always a part of the root system experience a mild water stress. Water was applied with 2 drip lines and 2 emitters per plant (0.50 m emitter spacing). Emitter flow was

$1.6 L h^{-1}$ (total flow: $3.2 L plant^{-1}$); [redacted] was applied to both sides of the root system. Water was applied with 1 drip line and 2 emitters per plant (0.50 m emitter spacing). Emitter flow was $1.6 L h^{-1}$ (total flow: $3.2 L plant^{-1}$). [redacted] non irrigated received only natural rainfall. There was only one soil column in the field study with PRD treatments achieved by applying different amounts of irrigation water to opposite sides of the plant. Air tem-

Field condition

Table 2


Effect of five irrigation treatments on stem diameter and height of papaya in a field study. July (2012) and October (2012).

Treatment ¹	Sampling Time			
	July		October	
	Diameter (mm)	Height (m)	Diameter (mm)	Height (m)
RDI	115.91	3.48ab ²	114.32	3.61
FI	110.52	3.31ab	115.35	3.41
NI-field	113.76	3.08c	113.87	3.65
PRD100	112.80	3.37ab	121.35	3.67
PRD70	117.85	3.57a	116.88	3.79
	ns ³		ns	ns

¹ FI=full irrigation; NI-field=no irrigation after treatment initiation; PRD100=partial root zone drying with 100% water replacement; PRD70=partial root zone drying with 70% water replacement; RDI=regulated deficit irrigation with 70% water replacement.

² Mean values within a column followed by the same letter are not significantly different ($P=0.05$) based on Tukey's multiple-range test.

³ Non-significant difference.






Table 4


Effect of 5 irrigation treatments on photosynthesis (A), stomatal conductance (G_s) and transpiration (E) of papaya in a field study, July (2012) and October (2012).

Treatment ¹	Sampling	A ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	G_s ($\text{mol m}^{-2} \text{s}^{-1}$)	E ($\text{mmol m}^{-2} \text{s}^{-1}$)
RDI 70%	July	14.8	0.19	4.3
FI	July	12.1	0.15	3.5
NI-field	July	13.2	0.21	4.4
PRD100	July	12.5	0.15	3.3
PRD70	July	12.7	0.18	4.0
		ns ²	ns	Ns
RDI 70%	October	9.5ab ²	0.12ab	4.4ab
FI	October	10.8a	0.13a	5.5a
NI-field	October	6.5b	0.06c	3.2b
PRD100	October	9.3ab	0.11abc	4.6ab
PRD70	October	7.2b	0.08bc	3.4b

¹ FI=full irrigation; NI-field=no irrigation after treatment initiation; PRD100=partial root zone drying with 100% water replacement; PRD70=partial root zone drying with 70% water replacement; RDI=regulated deficit irrigation with 70% water replacement.

² Mean values within a column followed by the same letter are not significantly different ($P=0.05$) based on Tukey's multiple range test.

³ Non-significant difference.






Fig. 2. Meteorological variables in a field study.

Table 3
Effect of 5 irrigation treatments on yield components and agronomic water use efficiency (AWUE) of papaya in a field study. March (2012) and June (2012).

Harvest	Treatment ^a	Number fruit plant ⁻¹	Average weight (g FW fruit ⁻¹)	Yield (kg FW ha ⁻¹)	kg FW plant ⁻¹	Irrigation + precipitation (L)	AWUE (kg FW fruit L ⁻¹)	AWUE (number fruit L ⁻¹)
March	FI	30b ^c	409ab	22,065ab	11.9b	2698	0.0044c	0.011c
March	NI-field	33ab	391b	23,991b	13.0ab	1025	0.0126a ←	0.032a ←
March	PRD100	38ab	436ab	31,290ab	16.9ab	2698	0.0063bc	0.014bc
March	PRD70	41a ←	437a ←	33,827a ←	18.3a ←	2189	0.0084b	0.019b
March	RDI	39ab	400ab	31,117ab	16.8ab	2189	0.0077b	0.018b
P=0.10								
June	FI	21a	244ab	9620ab	5.2ab	3755	0.0014	0.006
June	NI-field	6b ←	191b ←	2651b ←	1.4b ←	1107 ←	0.0013	0.006
June	PRD100	26a	309a	14,881	8.0a	3755	0.0021	0.007 ←
June	PRD70	19a	286a	10,390a	5.6ab	2949	0.0019	0.006
June	RDI	20a	282a	11,145a	6.0ab	2949	0.0020 ←	0.007
ns ^b								

^a FI=full irrigation; NI-field=no irrigation after treatment initiation; PRD100=partial root zone drying with 100% water replacement; PRD70=partial root zone drying with 70% water replacement; RDI=regulated deficit irrigation with 70% water replacement.

^b Mean values within a column followed by the same letter are not significantly different ($P=0.05$) based on Tukey's multiple range test.

^c Non-significant difference.

5. Conclusion

While there was evidence of non-hydraulic signals inducing stomatal closure in the PRD treatments compared to RDI in greenhouse studies, these effects were insufficient to alter dry matter partitioning, biomass, or yield components since there were no significant differences between PRD and RDI at either a 30% or 50% water deficit. A 50% water deficit in the greenhouse study for the PRD and RDI treatments was sufficient to significantly reduce biomass and dry matter partitioning compared to the FI treatment. In the field study, a 30% water deficit in both PRD70 and RDI treatments did not significantly reduce vegetative growth or yield components, compared to FI. It appears that papaya can tolerate moderate water deficits without a significant reduction in yield components indicating that <100% ET irrigation replacement may be scheduled but there is little or no difference between PRD and RDI. Further research will be needed to verify that moderate soil water deficits do not reduce quality.

Split root model - partial root volume irrigation

Minimal or no influence on:

- Relative water content of leaves
- Leaf expansion rate
- Stomatal conductance
- Net CO₂ assimilation
- Daily water use - gravimetric





One

Three

Two

Microsprinkler : 90°, 180°, or 360° application coverage

No influence on:


- Growth - height or stem diameter
- Date of first flowers
- Height of first fruit
- Yield

Papaya attributes that allow for this:


- 1. Efficient lateral transfer of water in stem**
- 2. Rapid root proliferation in wet zones**
- 3. Hydraulic redistribution into dry zones**

1. Efficient lateral water transfer in stem






20 cm above-
100%




10 cm above-
80%



6 cm above-
50%




Papaya attributes that allow for this:
2. Rapid root proliferation in wet
zones



-Trench profile


-Cores

Drip line locations



3. Hydraulic redistribution into dry zones

Midday transpiration



4 cm below
junction



Two hour chase period

Water Transfer in a Papaya-Corn Culture System

T.E. Marler
Western Pacific Tropical Research Center
University of Guam, Mangilao
Guam 96923
USA

Proc. Third IS on Papaya
Eds.: N. Chomchalow et al.
Acta Hort. 1022, ISHS 2014

Abstract

'Tainung 2' and 'Sunrise' papaya seedlings were grown in split-root containers. 'Honey Jean 3' sweet corn seeds were planted in one of the two containers that comprised each split-root papaya system. Following establishment of the corn seedlings, the papaya-corn systems were subjected to one of three treatments: 1) both halves of the papaya roots were well-watered (control); 2) both halves of the papaya roots received no water; (3) the papaya root half without the corn seedling was watered but the half with the corn seedling received no water. Pre-dawn leaf relative water content (RWC) and mid-morning stomatal conductance of corn leaves were the response variables used to quantify drought stress. Stomatal conductance reached zero by day 10, when RWC of treatment 2 plants was less than 50% and that of treatment 3 plants was 80%. At this stage, half of the remaining replications in treatment 3 were treated by cutting the connection between the roots in the dry compartment and the base of the papaya stem. This procedure relieved competition between the two species, but also eliminated the watered half of the papaya roots as a possible source of water for the corn plants. Leaf RWC of the corn plants relieved of papaya root competition declined to below that of corn plants within intact treatment 3 papaya split root systems. These results indicate hydraulic redistribution occurred from papaya roots in the watered pots to the corn plants. Water redistribution within papaya plants may have impacts on hydrologic processes, and should be considered when scaling fluxes to the orchard level.

Fl. We hypothesized that the difference observed in physiological response to PRD and RDI treatments between the papaya grown in the greenhouse and field may be related to the different volumes of soil explored by the root system. The physiological response of papaya to PRD and RDI was more affected in greenhouse-grown than field-grown papaya because in the greenhouse study, the roots are limited to the volume of the pot. In addition, in field conditions rainfall can increase water availability in the soil and the roots have a greater volume of soil to explore. Thus, environmental variables such as VPD and PAR can more severely affect the gas exchange and growth of plants grown in the greenhouse than plants grown under field condition. In addition, PRD and RDI can increase stomatal sensitivity to VPD ([Collins et al., 2010](#)).

Collins, M.J., Fuentes, S., Barlow, W.R., 2010. [Partial rootzone drying and deficit irrigation increase stomatal sensitivity to vapour pressure deficit in anisohydric grapevines. *Funct. Plant Biol.* 37, 128–138.](#)

Flooding

Papaya is considered a species sensitive to low oxygen availability in the soil (hypoxia), which is commonly caused by waterlogging (Ogden et al., 1981; Malo and Campbell, 1986)

Reduced oxygen can occur as a result of tropical storms that saturate the soil for several days, flood irrigation, as well as micro-irrigation practices that create microenvironments of reduced soil oxygen

A completely flooded soil can cause **death to papaya plants in 2 d** (Wolf and Lynch, 1940; Khondaker and Ozawa, 2007) **or 3 to 4 d** (Samson, 1980)




Figure 11 The time course of stomatal conductance of papaya leaves relative to that of control plants (8.54 mg/liter) as influenced by dissolved oxygen concentration in the nutrient solution. Data were standardized as a fraction of control. (Marier, T. E., unpublished data, 1990.)





2

11











Khondaker and Ozawa (2007) constructed chambers that controlled soil gas composition at ambient (20%), 18% and 11% oxygen; under soil oxygen at and below 18%, A, chlorophyll content, large and small roots, and shoot dry matter were all decreased

Papaya Plant Growth as Affected by Soil Air Oxygen Deficiency



N.A. Khondaker* and K. Ozawa
Okinawa Subtropical Station
Japan International Research Center for Agricultural Sciences (JIRCAS)
Ishigaki-shi, Okinawa 907-0002
Japan
Present address: Bangladesh Agricultural Research Council Dhaka 1215
Bangladesh



Proc. 1st IS on Papaya
Eds. M.E. Chinn and R.E. Paull
IS 2007





Box 1: 20% O₂
Box 2: 18% O₂
Box 3: 11% O₂





Papaya, considered sensitive to hypoxia, responds with **accentuated senescence (chlorotic leaves), leaf fall and does not recover after hypoxic conditions are removed** (Marler et al., 1994).

These studies indicate that papaya **is sensitive to small reductions in soil oxygen content** and it is likely that micro-irrigation saturation of a small portion of the soil is having some negative effects. Consequently, a welldrained soil is essential for high productivity.



	100% of roots submerged	
	Dissolved O ₂ concentration (mg l ⁻¹)	
Treatment	Day 1	Day 2
0 g CaO ₂	3.63 ± 0.92b	4.38 ± 0.89b
2.28 g CaO ₂	7.00 ± 0.76a	5.38 ± 1.70ba
4.57 g CaO ₂	8.03 ± 1.09a	7.19 ± 1.50a

et al., 2009a). Hydrogen peroxide decomposes in the soil, releasing O₂ which is needed for aerobic metabolism in the roots (Gil et al., 2009a,b). When H₂O₂ comes in contact with water, it reacts to give off 0.5 mol of O₂ per mole H₂O₂ as shown in the equation $\text{H}_2\text{O}_2 + \text{H}_2\text{O} \rightarrow 0.5\text{O}_2 + 2\text{H}_2\text{O}$ (Gil et al., 2009a). In soil, solid oxygen compounds (i.e., CaO₂, MgO₂) breakdown to H₂O₂ which then provides oxygen to the rhizosphere (Liu and Porterfield, 2014).